

Pesticide dependence and associated risks in Uruguayan agriculture: limitations in its approach

Dependencia y riesgos del uso de fitosanitarios en la agricultura uruguaya: limitaciones para su análisis

Dependência e riscos do uso de produtos fitossanitários na agricultura uruguaia: limitações à sua análise

Palladino, Cintia; García, Isabel; Fernández, Grisel

 **Cintia Palladino**

mariacintiapalladino@gmail.com
Universidad de la República, Uruguay

 **Isabel García**

isabelgarciaquirry@gmail.com
Universidad de la República, Uruguay

 **Grisel Fernández**

griselfe@gmail.com
Universidad de la República, Uruguay

Agrociencia Uruguay

Universidad de la República, Uruguay
ISSN-e: 2730-5066
Periodicity: Bilingual
vol. 27, e1156, 2023
agrociencia@fagro.edu.uy

Received: 03 January 2023

Accepted: 12 May 2023

Published: 23 May 2023

URL: <http://portal.amelica.org/ameli/journal/506/5063857009/>

DOI: <https://doi.org/10.31285/AGRO.27.1156>

Corresponding author: mariacintiapalladino@gmail.com

Agrociencia Uruguay, 2023



This work is licensed under Creative Commons Attribution 4.0 International.

Abstract: Pesticides have become an essential input for agriculture in the last decades. However, the growing concern about the potential impact produced by such dependency on human health and the environmental level has led to strong questionings about the use of pesticides worldwide. This paper aims to analyze the use of pesticides in Uruguay's agriculture and their potential risks with an emphasis on ecotoxicological risks. For such purposes, data on imports and use of pesticides in Uruguay and some other agricultural regions corresponding to the last two decades (2000-2020) were collected. In addition, information on the risks associated with the use of different pesticides compiled in the Pesticide Properties Database of the University of Hertfordshire was reviewed, as well as information generated by related national and international research. The information collected indicates high intensity usage of pesticides in Uruguayan agriculture and uncertainties in relation to environmental risks that may arise from the current forms of use.

Keywords: ecotoxicology, toxicology, environmental fate.

Resumen: Los fitosanitarios se han convertido en un insumo indispensable para la agricultura en las últimas décadas. Sin embargo, la creciente preocupación sobre los impactos que pueda generar esta dependencia a nivel de la salud humana y/o la sostenibilidad ambiental ha determinado fuertes cuestionamientos al uso de pesticidas a nivel mundial. El presente trabajo pretende analizar el uso de fitosanitarios en la agricultura de Uruguay y sus riesgos potenciales con énfasis en los ecotoxicológicos. Con tal objetivo, se procedió a la recopilación de información de datos de importación y uso de fitosanitarios a nivel nacional e internacional de otras regiones agrícolas, correspondientes a las dos últimas décadas (2000-2020). Complementariamente, se revisó la información de los riesgos asociados al uso de distintos pesticidas compilada en la base de datos de la Pesticide Properties Database de la Universidad de Hertfordshire, así como la generada en investigaciones nacionales e internacionales relacionadas con la temática. La información relevada señala alta intensidad del uso de fitosanitarios en la agricultura del país e incertidumbres en

relación con los riesgos ambientales y humanos que puedan derivarse de las formas de uso actual.

Palabras clave: ecotoxicología, toxicología, destino ambiental.

Resumo: Nas últimas décadas, os pesticidas tornaram-se um insumo indispensável para a agricultura. No entanto, a crescente preocupação com os impactos desta dependência na saúde humana e/ou na sustentabilidade ambiental levou a fortes questões sobre a utilização de pesticidas em todo o mundo. Este documento visa analisar a utilização de pesticidas na agricultura uruguaia e os seus riscos potenciais. Para tais fins, foram compiladas informações sobre a importação e utilização de produtos fitossanitários a nível nacional e internacional de outras regiões agrícolas durante as últimas duas décadas (2000-2020). Além disso, foram revistas informações sobre os riscos associados à utilização de diferentes pesticidas compiladas na Base de Dados de Propriedades dos Pesticidas da Universidade de Hertfordshire, bem como informações geradas na investigação nacional e internacional relacionada com o assunto. A informação coletada indica uma elevada intensidade de utilização de pesticidas na agricultura do país e incertezas em relação aos riscos ambientais e humanos que podem surgir das atuais formas de utilização.

Palavras-chave: ecotoxicologia, toxicologia, destino ambiental.

1. INTRODUCTION

Pesticides are one of the most important inputs in agricultural production. They effectively reduce losses caused by insects, diseases and weeds, improving crop yields and their quality⁽¹⁾⁽²⁾⁽³⁾. However, it has been widely demonstrated that, by reaching soil, water and air compartments, they also represent a major threat to environmental sustainability. Non-target organisms such as birds, fish, beneficial insects, plants and others are systematically exposed to pesticides and therefore affected by their toxicity⁽⁴⁾. Pesticides also pose serious risks to human health. Human exposure can occur directly due to occupational, agricultural, and household use, or indirectly, mainly through contaminated food and water consumption⁽⁵⁾⁽⁶⁾.

Despite the cumulative evidence generated in the past years and the growing awareness of negative impacts from pesticides; various attempts proposed to mitigate the overuse of pesticides have been practically ineffective. One of the most cited examples is the adoption of integrated pest management (IPM). Moss⁽⁷⁾ pointed out that IPM implementation has been poor, with little evidence of concomitant reductions in pesticide use.

Moreover, the quantity of pesticide production and consumption has kept increasing annually at the global level. According to the latest data from the Food and Agricultural Organization⁽⁸⁾, the total pesticide consumption worldwide has risen from 2,047,087 tons in 2000 to 2,611,124 tons in 2020.

Pesticide dependency is a fundamental part of modern agricultural regime and a very complex phenomenon. Even when pesticide resistance has been widely considered as one of the major causes⁽⁹⁾⁽¹⁰⁾. Hu⁽¹¹⁾ argues that pesticide dependency is not only a technological issue as pesticide resistance thesis

AUTHOR NOTES

mariacintiapalladino@gmail.com

indicates, but rather a man-made issue, with socioeconomic and political reasons for both the country and the farmers, involving multi-dimensions, multi-actors and multi-scales factors.

Consequently, multiple and complex actions at different levels are required when addressing pesticide dependency mitigation⁽¹²⁾. The actions to be developed in each region or country, as well as their prioritization and urgency, require previous and basic analysis of the intensity of pesticide use, the expected behavior in environmental fate and the associated risks⁽¹³⁾.

The objective of this work was to analyze pesticide use in Uruguay, combined with the information available on its potential risks with an emphasis on ecological risks.

2. MATERIALS AND METHODS

The information made available by the General Directorate for Agricultural Services (DGSA) from the Ministry of Livestock, Agriculture and Fisheries of Uruguay (MGAP)⁽¹⁴⁾, and the Food and Agriculture Organization of the United Nations (FAO)⁽⁸⁾ corresponding to the last two decades (2000-2022) was used for the diagnosis of pesticide use in Uruguay and other agricultures regions.

For the estimation of the most used active ingredients in the country, a sample was taken using information collected from the DGSA's system of pesticide application registrations from 2019 to 2022⁽¹⁵⁾. For the potential risks associated with its use, the information reviewed considered the Pesticide Properties Database⁽¹⁶⁾, national and international research related to the subject.

3. RESULTS AND DISCUSSION

3.1 Use of pesticides in Uruguayan agriculture

The use of these products has had a sustained increase since the year 2000, showing very similar trends to those observed worldwide. The total amount of active substances imported varied from 3.9 to 10.6 tons between 2000 and 2021⁽¹⁴⁾ (Figure 1).




FIGURE 1

Pesticide imports from 2000 to 2021 in Uruguay⁽¹⁴⁾ Values include raw material

Herbicides were the group that registered the greatest change, representing 89% of the total volume increase in kg or l of the formulates in 2021⁽¹⁴⁾. While comparing the worldwide use of pesticides in 2020, 53% is the share of herbicides, 18% is the share of insecticides, 23% is that of fungicides, and others account for 7% only⁽⁸⁾.

It should be clarified that pesticide products imports are not necessarily a synonymous of the use of such product. There is no information on how much of the imported pesticides are truly used in the country, nor if the use is exclusively for agricultural purposes. Despite this, it is an indicator commonly used for global and comparative analyses in this subject⁽¹⁷⁾.

The greater use of pesticide products in the country has a strong association with the agricultural area's growth in this period. Favourable agricultural prices and a rapid generalization of the practice of zero tillage were determinants of agriculture expansion towards non-traditional areas and mainly agriculture intensification process, as a result of massive adoption of double cropping⁽¹⁸⁾.

In this way, the increase in pesticide products imports reflects an increase in the intensity of their use, and therefore refers to greater quantities of pesticide products per hectare cultivated and per year. The information collected shows that higher amounts of pesticides per hectare in the periods between 2000 and 2020 varied from 2.58 to 7.97 kg/ha in Uruguay, just as it can be observed in Figure 2⁽⁸⁾. There are also marked differences between regions. While in Europe there is a tendency to decrease the intensity of pesticide use, the Southern Cone of South America shows an increasing trend, which is noticeable in the case of Uruguay.




FIGURE 2

Pesticides intensity use in Uruguay vs South American and European countries⁽⁸⁾

When we analyze the active ingredients comprising the groups of pesticide products imported in 2021⁽¹⁴⁾ in detail, a clear predominance of a few active ingredients was revealed (Table 1). The most notorious case is that of herbicides, where 88% corresponds to only 5, while 68 active ingredients correspond to the remaining 12%. In the case of imported fungicides and insecticides, this is not so evident, although a few active ingredients still predominate.

TABLE 1
The most imported active ingredients in Uruguay in 2021¹⁴

	Active ingredients	%
Herbicides	Glyphosate	67
	2,4-D	9
	S-metolachlor	6
	Clethodim	2
	Paraquat	3
	Others (68)	12
Insecticides	Chlorpyrifos	47
	Paraffin oil	12
	Triflumuron	6
	Aluminium phosphide	5
	Chlorantraniliprole	3
	Others (33)	27
Fungicides	Sodium metabisulphite	18
	Mancozeb*	18
	Copper oxide	14
	Ziram	8
	Captan	8
	Others (44)	35

* Mancozeb is also imported in mixtures with metalaxyl. These quantities were not considered because they are not separately reported.

Nevertheless, considering that pesticides are used at very different doses per hectare, kilograms of active substances do not reflect directly the use.

In this context of increasing pesticides imports, but mainly of higher intensity of use, it seems likely that associated risks would also increase, depending on their toxicological and ecotoxicological hazards.

3.2 Toxicology and ecotoxicology characterization of major used pesticides

In an attempt to make a rough estimate of the potential risks of the most commonly used active ingredients in Uruguay, a hazard profile was compiled (Table 2) using information from the DGSA's Register of Application of Pesticide⁽¹⁵⁾. It is important to clarify that this hazard profile only considered information on the parameters currently required to register pesticide products in Uruguay⁽¹⁹⁾.

TABLE 2
Toxicological profile according to Pesticide Properties Database⁽¹⁶⁾ of the main used pesticides (2019-2022)⁽¹⁵⁾

		Human health	Terrestrial ecotoxicology			Aquatic ecotoxicology	
		Mammals	Earthworms	Honeybees	Birds	Fish	Aquatic invertebrates
Insecticides	Chlorantraniliprole	Green	Green	Yellow	Green	Yellow	Red
	Triflumuron	Green	Yellow	Green	Yellow	Red	Red
	Chlorpyrifos	Red	Yellow	Red	Yellow	Red	Red
	Lambda-cyhalothrin	Red	Yellow	Red	Green	Red	Red
	Bifenthrin	Red	Yellow	Red	Yellow	Red	Red
Herbicides	Glyphosate	Green	Green	Green	Green	Green	Green
	2,4 D	Yellow	Yellow	Green	Yellow	Yellow	Green
	Clethodim	Green	Yellow	Yellow	Yellow	Yellow	Yellow
	Clopyralid	Green	Yellow	Yellow	Yellow	Yellow	Yellow
	S-metolachlor	Green	Yellow	Green	Yellow	Yellow	Yellow
Fungicides	Azoxystrobin	Green	Yellow	Green	Green	Yellow	Yellow
	Prothioconazole	Green	Yellow	Green	Green	Yellow	Yellow
	Pyraclostrobin	Green	Yellow	Green	Green	Red	Red
	Epoxiconazole	Green	Yellow	Green	Green	Yellow	Yellow
	Tebuconazole	Yellow	Green	Green	Yellow	Yellow	Yellow

Toxicity scale (Ts): mammals (rat) acute oral LD₅₀ (mg/kg) < 5-50 = very hazardous (red); 50-2000 = moderately hazardous (yellow); > 2000 = unlikely to present an acute hazard (green). Ts in earthworms (*Eisenia foetida*) acute 14-day LC₅₀ (mg/kg), > 1000 = Low 10 - 1000 = Moderate < 10 = High. Ts in birds (*Oncorhynchus mykiss*/ *Colinus virginianus*) Acute LD₅₀ (mg/kg) > 2000 = low (green); 100 - 2000 = Moderate (orange); < 100 = High (red). Ts in honeybees (*Apis mellifera*) acute LD₅₀ (µg/bee) > 100 = Low (green); 1 - 100 = Moderate (orange); < 1 = High (red). Ts in fish (*Colinus virginianus*) acute 96 h LD₅₀ (mg/l) > 100 = Low (green); 0,1 - 100 = Moderate (orange); < 0,1 = High (red); Aquatic invertebrates (*Daphnia magna*) acute 48 hour EC₅₀ (mg/l) > 100 = Low; 0,1 - 100 = Moderate and < 0,1 = High.

The danger for humans from pesticide products is classified into toxicological categories. These categories are defined according to the acute risk in mammals, resulting from a single or repeated exposure over a relatively short period of time with the pesticide product. Based on this classification, we notice that in the case of herbicides and fungicides predominant substances do not represent a major threat to human health (Table 2). However, in the case of insecticides, those which predominate are highly dangerous to humans.

National epidemiology studies in human toxicology have identified a wide range of conditions and diseases associated with environmental exposure to pesticides. Evidence has been found in the working environment, in rural and urban populations associated with agriculture⁽²⁰⁾⁽²¹⁾⁽²²⁾; and in the general unexposed population⁽²³⁾.

The human health risk assessment is complex due to its strong dependence on periods and levels of exposure, type of pesticides used, and characteristics of the environment and the human communities of the areas where pesticides are applied. Long-term, standardized and validated studies are needed to confirm the link between pesticides and their negative effects on human health. In addition, the risk to human health should be assessed not only for specific active substances and different formulations, but also for the possible cumulative and interactive effects of exposure to multiple pesticides over time⁽²⁴⁾⁽²⁵⁾. It is firmly suggested that risk assessment of pesticide-exposed populations use occupational exposure matrices that consider the tasks performed, the crops and active ingredients, and more complete occupational histories, which could reduce errors due to exposure⁽²⁶⁾.

Regarding the ecotoxicological characteristics of the profile (Table 2), the main insecticides used present more toxicity in terrestrial and aerial organisms in relation to the other groups. However, in the case of aquatic organisms, all three groups of pesticide products present toxicity problems.

Ecotoxicological studies are also complex. They are performed with standard organisms generally selected according to the simplicity of their laboratory study. These organisms are not necessarily the organisms of interest in all the environments in which the pesticide product would be applied⁽²⁷⁾.

In addition, much of the information currently included in the pesticide registry is based on toxicity studies under laboratory conditions, focusing on groups of individual organisms and evaluating each active ingredient separately. Although these types of tests allow methodological homogenization and easily comparable results, they may be questionable in their ability to effectively protect ecosystems, given the differences between the assessment conditions and the conditions of real use⁽²⁸⁾⁽²⁹⁾.

Currently, and especially in extensive agriculture, pesticides are used in mixtures, and may generate synergistic toxicological effects, greater than those tested individually. In addition, these systems already have pesticide products in the process of degradation, generating cocktails of numerous compound that lead to stressful conditions for non-targeted organisms. In addition, risk assessments assess the effects on individual organisms, but not on population or community level⁽³⁰⁾.

Also, information regarding the potential effects of the resulting secondary metabolites from degradation is very limited. They have their own toxicological and ecotoxicological characteristics and can be even more toxic than the parent compounds⁽³¹⁾.

Moreover, the risk generated by a compound is not only linked to the dose and metabolites it generates, but is also dependant on its environmental behaviour. A clear example is the persistence or active bioavailability of pesticide residues over time in different environmental compartments, which is strongly dependent on

the physio-chemical characteristics of the compound and the variable characteristics of the destination compartment⁽¹⁶⁾⁽³²⁾⁽³³⁾. In this sense, glyphosate is considered of low risk in soil, among other aspects based on its categorization as non-persistent. At present, it is known that it may persist according to its form of use, to the point that its reclassification as pseudo-persistent is recommended⁽³⁴⁾.

Pesticide residues cause direct and indirect negative effects on non-targeted organisms⁽²³⁾, microorganisms, flora, physicochemical, and biological properties of agricultural soil⁽³⁵⁾⁽³⁶⁾.

In Uruguay, the presence of pesticide residues has been reported in watercourses and drinking water⁽³⁷⁾⁽³⁸⁾, in soil⁽³⁹⁾⁽⁴⁰⁾, in the air⁽⁴¹⁾, in beehives and in component parts of beehives⁽⁴²⁾⁽⁴³⁾⁽⁴⁴⁾⁽⁴⁵⁾, and in agrifoods⁽⁴⁶⁾⁽⁴⁷⁾⁽⁴⁸⁾. Furthermore, there are several studies on the effect of the use of pesticide products at the biota level in environmental compartments⁽⁴⁹⁾⁽⁵⁰⁾⁽⁵¹⁾⁽⁵²⁾⁽⁵³⁾⁽⁵⁴⁾⁽⁵⁵⁾⁽⁵⁶⁾⁽⁵⁷⁾.

Based on the toxicological-ecotoxicological parameters and environmental fate of an active ingredient, pesticide products are registered, suspended, restricted or prohibited for their use.

There are currently 2601 pesticide products registered by MGAP in Uruguay. Eighty one of the registered active ingredients (which comprise 330 products containing these active ingredients) are classified as highly hazardous pesticides (HHPs)⁽⁵⁸⁾. HHPs are defined by the World Health Organisation, the Globally Harmonised System of Classification and Labelling of Chemicals, Pesticide Action Network International and FAO⁽⁵⁹⁾⁽⁶⁰⁾ as pesticides that pose particularly high acute or chronic risks to human health or the environment. The European Union and many other countries have banned 41 of the 81 active ingredients that are classified as HHPs and that are registered and commercially available in Uruguay⁽⁵⁸⁾.

Uruguay has only 22 banned active ingredients according to PAN⁽⁶¹⁾, similar to Argentina, Chile and Paraguay (18, 27 and 11 active ingredients, respectively), while Brazil has 133 banned active ingredients. Countries from the European Union are more restrictive with the use and registration of pesticide products⁽⁶¹⁾. When we relate this information with that corresponding to the effectively used amounts, we notice that the countries with the greatest use of pesticide products are those with the least restrictions.

It is worth mentioning that in countries with more restrictions the environmental risk assessment required for the registration of new pesticide products must be carried out in different cultivation scenarios, soil and climate representative of the region in which they will be used⁽²⁴⁾. Despite the associated costs of these types of studies, they are essential to better estimate the impact of pesticide use on a particular ecosystem.

4. FINAL CONSIDERATIONS

The available information indicates a high intensity of pesticide use in Uruguayan agriculture. It also evidences the presence of pesticide residues in different environmental compartments, and several negative effects have been reported on local non-target organisms.

Improvements on the registration and renewal system of pesticides, as adopted by countries with more restricted regulations, could be an effective strategy to mitigate deleterious effects of pesticides use.

REFERENCES

1. Fenik J, Tankiewicz M, Biziuk M. Properties and determination of pesticides in fruits and vegetables. *TrAC Trends Anal Chem.* 2011;30:814-26.
2. Strassemeyer J, Daehmlow D, Dominic A, Lorenz S, Golla B. SYNOPSIS-WEB, an online tool for environmental risk assessment to evaluate pesticide strategies on field level. *Crop Prot.* 2017;97:28-44.

3. Keulemans W, Bylemans D, De Coninck B. Farming without plant protection products: Can we grow without using herbicides, fungicides and insecticides? [place unknown]: European Parliamentary Research Service; 2019. 44p. Doi: 10.2861/05433.
4. Sharma A, Shukla A, Attri K, Kumar M, Kumar P, Suttee A, Singh G, Barnwal RP, Singla N. Global trends in pesticides: a looming threat and viable alternatives. *Ecotoxicol Environ Saf.* 2020;201:110812. Doi: 10.1016/j.ecoenv.2020.110812.
5. Li Z, Jennings A. Worldwide regulations of standard values of pesticides for human health risk control: a review. *Int J Environ Res Public Health.* 2017;14(7):826. Doi: 10.3390/ijerph14070826.
6. Rani L, Thapa K, Kanojia N, Sharma N, Singh S, Grewal AS, Srivastav AL, Kaushal J. An extensive review on the consequences of chemical pesticides on human health and environment. *J Clean Prod.* 2020;283:124657. Doi: 10.1016/j.jclepro.2020.124657.
7. Moss S. Integrated weed management (IWM): why are farmers reluctant to adopt non-chemical alternatives to herbicides? *Pest Manag Sci.* 2019;75(5):1205-11.
8. FAO. FAOSTAT [Internet]. [place unknown]: FAO; 2023 [cited 2023 May 15]. Available from: <http://www.fao.org/faostat/es/#data/FBS>
9. Oerke E. Crop losses to pests. *J Agric Sci.* 2006;144(1):31-43.
10. Bakker L, Van Der Werf W, Titttonell P, Wyckhuys KAG, Bianchi FJJA. Neonicotinoids in global agriculture: evidence for a new pesticide treadmill? *Ecol Soc.* 2020;25:26. Doi: 10.5751/ES-11814-250326.
11. Hu Z. What socio-economic and political factors lead to global pesticide dependence? a critical review from a social science perspective. *Int J Environ Res Public Health.* 2020;17(21):8119. Doi: 10.3390/ijerph17218119.
12. Ríos-González A, Jansen K, Sánchez-Pérez HJ. Pesticide risk perceptions and the differences between farmers and extensionists: towards a knowledge-in-context model. *Environ Res.* 2013;124:43-53.
13. Zhang Z, Yan X, Jones K, Jiao C, Liu Y, Zhu Y, Zhang Q, Zhai L, Shen Z, Chen L. Pesticide risk constraints to achieving Sustainable Development Goals in China based on national modeling. *NPJ Clean Water.* 2022;5:59. Doi: 10.1038/s41545-022-00202-0.
14. Ministerio de Ganadería, Agricultura y Pesca, DGSA (UY). Consulta de productos fitosanitarios [Internet]. Montevideo: MGAP; 2022 [cited 2023 May 15]. Available from: <https://www.mgap.gub.uy/profit/pantallaweb.aspx>
15. República Oriental del Uruguay, Ministerio de Ganadería, Agricultura y Pesca. Requisitos para la inscripción en el Registro Único de Operadores (RUO). Resolución N° 959/022 [Internet]. 2023 [cited 2023 May 15]. Available from: <https://bit.ly/42ALPpn>
16. PPDB: Pesticide Properties Database [Internet]. Hatfield: University of Hertfordshire; 2020 [cited 2023 May 15]. Available from: <http://sitem.herts.ac.uk/aeru/ppdb/>
17. Pesticides use, pesticides trade and pesticides indicators: global, regional and country trends, 1990–2020. Rome: FAO; 2022. 13p. (FAOSTAT Analytical Brief; 46). Doi: 10.4060/cc0918en.
18. Figueredo S, Guibert M, Arbeletche P. Ciclo sojero y estrategias de los actores de la producción agropecuaria en el litoral uruguayo. *Eutopía.* 2019;(16):99-110. Doi: 10.17141/eutopia.16.2019.4103.
19. Ministerio de Ganadería, Agricultura y Pesca, DGSA (UY). Registro y renovación de productos fitosanitarios (Plaguicidas Agrícolas) [Internet]. Montevideo: MGAP; 2023 [cited 2023 May 15]. Available from: <https://bit.ly/3ObbUXr>
20. Burger M, Pose Román D. Plaguicidas salud y ambiente: experiencia en Uruguay. Montevideo: Universidad de la República; 2012. 369p.
21. Heinzen J, Rodríguez N. Procesos destructores para la salud vinculados a la manipulación de agroquímicos en trabajadores agrícolas de Young, Uruguay. *Cien Trab.* 2016;18(56):117-23.
22. Rodríguez N, Heinzen J. Producciones desentido en torno a la exposición a agroquímicos: el caso de la ciudad de Young. *INNOTEC.* 2017;13:115-24.

23. Laborde A, Tomasina F, Bianchi F, Bruné MN, Buka I, Comba P, Corra L, Cori L, Duffert CM, Harari R, Iavarone I, McDiarmid MA, Gray KA, Sly PD, Soares A, Suk WA, Landrigan PJ. Children's health in Latin America: the influence of environmental exposures. *Environ Health Perspect.* 2015;123(3):201-9. Doi: 10.1289/ehp.1408292.
24. Ntzani EE, Chondrogiorgi M, Ntritsos G, Evangelou E, Tzoulaki I. Literature review on epidemiological studies linking exposure to pesticides and health effects. *EFSA Support Publ.* 2013;10(10):497E. Doi: 10.2903/sp.efsa.2013.EN-497.
25. Zúñiga-Venegas LA, Hyland C, Muñoz-Quezada MT, Quirós-Alcalá L, Butinof M, Buralli R, Cardenas A, Fernandez RA, Foerster C, Gouveia N, Gutiérrez Jara JP, Lucero BA, Muñoz MP, Ramírez-Santana M, Smith AR, Tirado N, van Wendel de Joode B, Calaf GM, Handal AJ, Soares da Silva A, Cortés S, Mora AM. Health effects of pesticide exposure in Latin American and the Caribbean Populations: a scoping review. *Environ Health Perspect.* 2022;130(9):96002. Doi: 10.1289/EHP9934.
26. Carles C, Bouvier G, Lebailly P, Baldi I. Use of job-exposure matrices to estimate occupational exposure to pesticides: a review. *J Expo Sci Environ Epidemiol.* 2017;27(2):125-40.
27. Rutgers M, Schouten AJ, Bloem J, Van Eekeren N, De Goede RGM, Jagersop Akkerhuis GAJM, Breure AM. Biological measurements in a nationwide soil monitoring network. *Eur J Soil Sci.* 2009;60(5):820-32.
28. Topping CJ, Aldrich A, Berny P. Overhaul environmental risk assessment for pesticides. *Science.* 2020;367(6476):360-3.
29. Bopp SK, Kienzler A, Richarz AN, van der Linden SC, Paini A, Parissis N, Worth AP. Regulatory assessment and risk management of chemical mixtures: challenges and ways forward. *Crit Rev Toxicol.* 2019;49(2):174-89.
30. Vryzas Z, Ramwell C, Sans C. Pesticide prioritization approaches and limitations in environmental monitoring studies: from Europe to Latin America and the Caribbean. *Environ Int.* 2020;143:105917. Doi: 10.1016/j.envint.2020.105917.
31. Michel C, Baran N, André L, Charron M, Joulain C. Side effects of pesticides and metabolites in groundwater: impact on denitrification. *Front Microbiol.* 2021;12:662727. Doi: 10.3389/fmicb.2021.662727.
32. Tariq SR, Nisar L. Reductive transformation of profenofos with nanoscale Fe/Ni particles. *Environ Monit Assess.* 2018;190(3):123. Doi: 10.1007/s10661-018-6500-2.
33. Wu L, Chládková B, Lechtenfeld OJ, Lian S, Schindelka J, Herrmann H, Richnow HH. Characterizing chemical transformation of organophosphorus compounds by ¹³C and ²H stable isotope analysis. *Sci Total Environ.* 2018;615:20-8. Doi: 10.1016/j.scitotenv.2017.09.233.
34. Primost JE, Marino DJG, Aparicio VC, Costa JL, Carriquiriborde P. Glyphosate and AMPA, "pseudo-persistent" pollutants under real-world agricultural management practices in the Mesopotamic Pampas agroecosystem, Argentina. *Environ Pollut.* 2017;229:771-9. Doi: 10.1016/j.envpol.2017.06.006.
35. Alengebawy A, Abdelkhalek ST, Qureshi SR, Wang MQ. Heavy metals and pesticides toxicity in agricultural soil and plants: ecological risks and human health implications. *Toxics.* 2021;9(3):42. Doi: 10.3390/toxics9030042.
36. Jacobsen CS, Hjelmsø MH. Agricultural soils, pesticides and microbial diversity. *Curr Opin Biotechnol.* 2014;27:15-20. Doi: 10.1016/j.copbio.2013.09.003.
37. Ernst F, Alonso B, Colazzo M, Pareja L, Cesio V, Pereira A, Pérez-Parada A. Occurrence of pesticide residues in fish from south American rainfed agroecosystems. *Sci Total Environ.* 2018;631-632:169-79. Doi: 10.1016/j.scitotenv.2018.02.320.
38. Planes de monitoreo: Informes Comité Científico: Subprograma 1: Monitoreo de la calidad de agua, sedimento y biota en el río Uruguay elaborados por el Comité Científico [Internet]. Paysandú: Caru; 2023 [cited 2023 May 15]. Available from: <https://www.caru.org.uy/web/actividades/planes-de-monitoreo/>
39. Colazzo M, Pareja L, Cesio V, Heinzen H. Multi-residue method for trace pesticide analysis in soils by LC-QQQ-MS/MS and its application to real samples. *Int J Environ Anal Chem.* 2018;98(14):1292-308.
40. Alonso B, Griffero L, Bentos Pereira H, Pareja L, Pérez Parada A. Determination of glyphosate and AMPA in freshwater and soil from agroecosystems by 9-fluorenylmethoxycarbonyl chloride derivatization and liquid

chromatography - fluorescence detection and tandem mass spectrometry. *MethodsX*. 2022;9:101730. Doi: 10.1016/j.mex.2022.101730.

41. Besil N, Hladki R, Rivero F, Cesio MV, Heinzen H. Pesticide residues in rainwater from the northwest region of Uruguay: method validation and seasonal analysis. In: 14th International IUPAC Congress on Crop Protection and 71st International Symposium on Crop Protection; 2019 May 19-24; Ghent, Belgium. [place unknown: publisher unknown]; 2019. p. 35.
42. Niell S, Jesús F, Pérez C, Mendoza Y, Díaz R, Franco J, Cesio V, Heinzen H. QuEChERS adaptability for the analysis of pesticide residues in beehive products seeking the development of an agroecosystem sustainability monitor. *J Agric Food Chem*. 2015;63(18):4484-92. Doi: 10.1021/acs.jafc.5b00795.
43. Murcia-Morales M, Heinzen H, Parrilla-Vázquez P, Gómez-Ramos MDM, Fernández-Alba AR. Presence and distribution of pesticides in apicultural products: a critical appraisal. *TrAC Trends Anal Chem*. 2022;146:116506. Doi: 10.1016/j.trac.2021.116506.
44. Pareja L, Colazzo M, Pérez-Parada A, Niell S, Carrasco-Letelier L, Besil N, Cesio MV, Heinzen H. Detection of pesticides in active and depopulated beehives in Uruguay. *Int J Environ Res Public Health*. 2011;8(10):3844-58. Doi: 10.3390/ijerph8103844.
45. Niell S, Jesús F, Pérez N, Pérez C, Pareja L, Abbate S, Carrasco-Letelier L, Díaz S, Mendoza Y, Cesio V, Heinzen H. Neonicotinoids transference from the field to the hive by honey bees: towards a pesticide residues biomonitor. *Sci Total Environ*. 2017;581-582:25-31.
46. Pareja L, Colazzo M, Pérez-Parada A, Besil N, Heinzen H, Böcking B, Cesio V, Fernández-Alba AR. Occurrence and distribution study of residues from pesticides applied under controlled conditions in the field during rice processing. *J Agric Food Chem*. 2012;60(18):4440-8.
47. Pareja L, Martínez-Bueno MJ, Cesio V, Heinzen H, Fernández-Alba AR. Trace analysis of pesticides in paddy field water by direct injection using liquid chromatography-quadrupole-linear ion trap-mass spectrometry. *J Chromatogr A*. 2011;1218(30):4790-8. Doi: 10.1016/j.chroma.2011.02.044.
48. Palladino C, Puigvert F, Muela A, Taborda B, Pérez CA, Pérez-Parada A, Pareja L. Evaluation of Fusarium mycotoxins and fungicide residues in barley grain produced in Uruguay. *J Agric Food Res*. 2021;3:100092. Doi: 10.1016/j.jafr.2020.100092.
49. Porrini MP, Garrido PM, Umpiérrez ML, Porrini LP, Cuniolo A, Davyt B, González A, Eguaras MJ, Rossini C. Effects of synthetic acaricides and nosema ceranae (Microsporidia: Nosematidae) on molecules associated with chemical communication and recognition in honey bees. *Vet Sci*. 2020;7(4):199. Doi: 10.3390/vetsci7040199.
50. Castelli L, Balbuena S, Branchiccela B, Zunino P, Liberti J, Engel P, Antúnez K. Impact of chronic exposure to sublethal doses of glyphosate on honey bee immunity, gut microbiota and infection by pathogens. *Microorganisms*. 2021;9(4):845. Doi: 10.3390/microorganisms9040845.
51. Jorge-Escudero G, Pérez Polanco M, Lagerlöf JE, Pérez CA, Míguez D. Commercial fungicide toxic effects on terrestrial non-target species might be underestimated when based solely on active ingredient toxicity and standard earthworm tests. *Toxics*. 2022;10(9):488. Doi: 10.3390/toxics10090488.
52. Balbuena S, Castelli L, Zunino P, Antúnez K. Effect of chronic exposure to sublethal doses of imidacloprid and nosema ceranae on immunity, gut microbiota, and survival of africanized honey bees. *Microb Ecol*. 2023;85(4):1485-97. Doi: 10.1007/s00248-022-02014-8.
53. Lacava M, García LF, Viera C, Michalko R. The pest-specific effects of glyphosate on functional response of a wolf spider. *Chemosphere*. 2021;262:127785. Doi: 10.1016/j.chemosphere.2020.127785.
54. Benamú MA, Schneider MI, González A, Sánchez NE. Short and long-term effects of three neurotoxic insecticides on biological and behavioral attributes of the orb-web spider *Alpaida veniliae* (Araneae, Araneidae): implications for IPM programs. *Ecotoxicology*. 2013;22(7):1155-64.
55. Benamú M, Lacava M, García LF, Santana M, Fang J, Wang X, Blamires SJ. Nanostructural and mechanical property changes to spider silk as a consequence of insecticide exposure. *Chemosphere*. 2017;181:241-9.
56. Bueno AF, Bueno RCOF, Parra JRP, Vieira SS. Effects of pesticides used in soybean crops to the egg parasitoid *Trichogramma pretiosum*. *Ciênc Rural*. 2008;38(6):1495-503.

57. Dake RB, Bhamare VK, Kapare RB. Effect of Bio-Rational insecticides on Lady Bird beetles of sunflower ecosystem. *Bull Env Pharmacol Life Sci*. 2017;6(2):74-6.
58. Cárcamo MI. Los Plaguicidas Altamente Peligrosos (PAP) en Uruguay [Internet]. [place unknown]: IPEN; 2020 [cited 2023 May 15]. 97p. Available from: <https://bit.ly/3oa9Qo1>
59. Highly hazardous pesticides [Internet]. [place unknown]: PAHO; 2019 [cited 2023 May 15]. Available from: <https://www.paho.org/en/topics/highly-hazardous-pesticides>
60. International Code of Conduct on Pesticide Management: guidelines on highly hazardous pesticides [Internet]. Rome: FAO; 2016 [cited 2023 May 15]. 28p. Available from: <https://bit.ly/3O9nHWc>
61. PAN International consolidated the list of banned pesticides [Internet]. [place unknown]: PAN; 2022 [cited 2023 May 15]. Available from: <https://pan-international.org/pan-international-consolidated-list-of-banned-pesticides/>

ALTERNATIVE LINK

<https://agrocienciauruguay.uy/index.php/agrociencia/article/view/1156/1389> (pdf)