



Efecto del estrés por inundaciones en cacao (*Theobroma cacao* L.)

Saravia-Castillo, Gabriela; Castro-Cepero, Viviana; Julca Otiniano, Alberto; Alvarado-Huamán, Leonel; Borjas-Ventura, Ricardo

- Saravia-Castillo, Gabriela
National Agrarian University La Molina, Perú
- Castro-Cepero, Viviana
National Agrarian University La Molina, Perú
- Julca Otiniano, Alberto
National Agrarian University La Molina, Perú
- Alvarado-Huamán, Leonel
National Agrarian University La Molina, México
- Borjas-Ventura, Ricardo *
rborjas@lamolina.edu.pe
National Agrarian University La Molina, Perú

Journal of the Selva Andina Biosphere

Selva Andina Research Society, Bolivia
ISSN: 2308-3867
Periodicity: Bilingual
vol. 10, no. 2, 2022
directoreditorbiosphere@gmail.com

Received: 01 March 2022
Corrected: 01 August 2022
Accepted: 01 September 2022
Published: 01 November 2022

URL: <http://portal.amelica.org/ameli/journal/71/713755005/>

DOI: <https://doi.org/10.36610/j.jsab.2022.100200078x>

Selva Andina Research Society



This work is licensed under Creative Commons Attribution-NonCommercial 4.0 International.

Abstract: Despite the fact that cocoa is very important for small farmers, information about flooding stress is scarce and dispersed being a serious problem. Flood affects cocoa trees in different levels. For example, this kind of stress disrupt the physiology in term of photosynthesis and stomatal conductance; likewise, it provokes damages in the photosystem II reason for which plants of cocoa under flood display high value of Fv/Fm. On the other hand, the overabundance of water reduces the capacity plant have to take nutrients (macro and micro nutrients) from soil. The Reduction of growth and the yield are other symptoms that cocoa trees may be under anoxic conditions. It was interesting to find that cocoa has biochemical strategy to overcome lack of oxygen such as increasing the glycolysis. This specie also has morphological strategy such as forming lenticels. It is very important to choose a flooding stress tolerant cultivar to limit the negative effects of this abiotic stress.

Keywords: Waterlogging, physiology, biochemistry, nutrition, production.

Resumen: El estrés por inundación en cacao es un problema cuya información está dispersa y es escasa, a pesar de que este cultivo es importante para los pequeños agricultores. Las inundaciones afectan a los árboles de cacao en diferentes niveles. Por ejemplo, este tipo de estrés altera la fisiología en términos de fotosíntesis y conductancia estomática; asimismo, provoca daños en el fotosistema II por lo que las plantas de cacao bajo inundación presentan valores elevados de Fv/Fm. Por otro lado, el exceso de agua reduce la capacidad que tienen las plantas de tomar nutrientes (macro y micronutrientes) del suelo. La reducción del crecimiento y el rendimiento son otros síntomas de que los árboles de cacao pueden estar en condiciones anóxicas. Fue interesante encontrar que el cacao tiene una estrategia bioquímica para superar la falta de oxígeno, como aumentar la glucólisis. También tiene una estrategia morfológica como formar lenticelas. Es muy importante elegir un cultivar tolerante al estrés por inundación para limitar los efectos negativos de estrés abiótico.

Palabras clave: Anegamiento, fisiología, bioquímica, nutrición, producción.

INTRODUCTION

The cocoa production is the main livelihood of 5-6 million of smallholders distributed in developing countries as Ivory Coast, Ghana, Indonesia, Nigeria, Cameroon, Brazil, Ecuador and Peru^{1,2}. Within those countries, the cocoa production is located in forestland where it has an important ecological role maintaining the biodiversity of insects and plants, and increasing the carbon sequestration³⁻⁶. Likewise, cocoa production has key social role since it generates employment. For example, in Peru, around of 190 thousand families and 450 thousand people both depend on cocoa crop⁷ (Figure 1).



FIGURE 1
Wild Cocoa tree planted in Lima, Peru

It is indubitable that crop productivity is associated to climatic conditions, hence any modification in this can disturb the development and growth of the cocoa trees. Therefore, there are two phenomena associated to climate change: drought and flooding. Although, the damage caused by water deficit is more intense than water excessive⁸, however, the waterlogging can disrupt the good performance of photosynthesis⁹ and it decreases the yield of crops¹⁰. For that reason, it can be considered a threat to the food security¹¹.

In the last years, the global climate has come changing dramatically, increasing among other parameters, the presence of heavy rain in short periods¹² putting the production of certain crops at risk, especially those grown on forestland or near to river bank as cocoa (*Theobroma cacao* L.). This is due to the fact that overabundance of water on soil decreases dramatically the presence of available O₂ (anoxic) to roots provoking physiological, morphological and biochemical changes of the cocoa plant, causing decreased yields^{13,14} (Figure 2).

AUTHOR NOTES

* Contact address: National Agrarian University La Molina. Faculty of Agronomy. Academic Department of Phytotechnics. Av. La Molina s/n. La Molina, Peru. Tel. +51 01614-7800.
Ricardo Borjas Ventura E-mail address: rborjas@lamolina.edu.pe




FIGURE 2

Effect of flooding in plant development. Prolonged periods of rain may provoke flooding that, at the same time, decreases the availability of O₂ to plants. In these circumstances the respiration is disrupted, and consequently the ATP synthesis. All those changes can negatively affect the physiology, nutrient uptake, growth and yield of crops, particularly cocoa

In order to face this abiotic problem, we need to know the impact of flooding on different aspects that determine the adequate growth and yield in cocoa trees. In spite of the waterlogging stress can potentially increase its negative impact on cocoa plantations, information on the issue is scarce. Additionally, the material relates to this stress is dispersed. In this context, this review aims to group and classify all information of the impact of flooding on the cocoa plant.

DEVELOPMENT

Effect on physiology of cocoa. Photosynthesis is one of the most important physiological processes in plants¹⁵ being responsible for synthesizing carbohydrates, that will be used to maintain the growth and development of crops¹⁶. To carry out this vital process the plants count on a photosynthetic apparatus that transforms the sunlight in energy.

That process is highly dependent on the entry of CO₂¹⁷. Therefore, whatever that disrupt the entry of CO₂ into plants can impair the photosynthates production. This relationship also denotes the high impact of abiotic factors, such as flood, on photosynthesis.

When the soil is totally covered by water, cocoa physiology in terms of photosynthesis and stomatal conductance^{9,18} are affected. In fact, the former can be reduced by 60-83 %^{13,19}, meanwhile, the latter can reduce by 81 %¹³. Furthermore, the transpiration is reduced by almost 60 %. The reduction in both photosynthesis rate and stomatal conductance could be associated to stomatal closure²⁰ that, at the same time, could be correlated to hormones such as ethylene²¹.

On the other hand, the photosynthetic apparatus is formed by Photosystems (FS) (I and II) whose functions are to catalyse the transfer the electrons from plastocyanin to ferredoxin (FS I) and to oxidize water (FS II)^{22,23}. One of the methods to measure impact of abiotic stress on PS II is quantifying the fluorescence²⁴. Interestingly, although some researchers as Bertolde et al.¹³ documented that Fv/Fm can be negatively affected by excess water in the soil, other such as Costa et al.²⁵ showed that photosynthetic efficiency (Fv/Fm) was almost constant in plant under flood stress, but the same plants rapidly modified the total performance index (PI_{tot}), suggesting the latter may be better parameter to evaluate the impact of waterlogging on plant vitality²⁶. According to Živčák et al.²⁷ and Swoczyna et al.²⁸, PI_{tot} is related to quantum efficiency to reduce the PSI

with photons from PSII. Regarding PSII fluorescence, da Silva Branco et al.²⁹ informed that flooding stress decreases q_p as tolerant cultivar as resistant cultivar. Nevertheless, cocoa plants under anoxic conditions can slightly raise the water use efficiency (WUE)³⁰.

Effect on biochemistry of cocoa. The activity of different protective enzymes allow plant to survive adverse conditions. The enzymatic response depends on the genotype as evidenced by Bertolde et al.¹³ in an experiment in which they observed that tolerant and susceptible genotypes to flooding showed differentiated enzymatic activity (Peroxidase and Polyphenol oxidases) compared to Control. The plant response also depends on time of exposure.

Furthermore, the content of starch and soluble sugars, in general, was reduced in the roots of plants subjected to waterlogging¹³, pointing out the high susceptibility of this organ to excessive water. In addition, proline might be an indicator of stress due to excessive water, but only for short term response, although it depends on cultivar studied⁸.

In this review, we found low information about the relation between anoxic (waterlogging) and respiration. However, the data found demonstrate that in flood-tolerant genotypes of cacao (TSA-792), the activity of alcohol dehydrogenase (EC 1.1.1.1), pyruvate carboxylase (EC 6.4.1.1) and lactate dehydrogenase increases, suggesting that an increment of glycolysis may be proper strategy used by tolerant genotypes¹³. This strategy is mentioned by other authors such as³¹ who also reported that glycolysis helps plants to produce rapidly ATP.

Effect on nutrient uptake. The plant nutrients are many important because ones prompt the growth and development of plants, Therefore, a deficiency in their content can affect the yield of plant as cocoa. The levels of nutrient in plants are severely affected by the excessive water on soil. Macronutrient such as N, P, K, Ca and Mg were negatively affected by overabundance of water on soil^{13,32}. In the case of micronutrients (Fe, Zn, Cu and Mn), the cocoa plants (both tolerant and susceptible to waterlogging) under flood stress had low absorption of these elements.

Effect on growth and morphology of cocoa. Inhibition of plant growth caused by waterlogging is reported in many crops, such as tomato³³ and oilseed rape³⁴. In the case of cocoa, the suppression of vegetative growth by flood stress is pointed out by de Almeida & Valle¹⁴ who stated that these phenomena limit the initial growth of cocoa seedlings diminishing the leave area and provoking the falling of leaves.

Likewise, in an experiment with a four months old seedling subjected to waterlogging by one month, the researchers noticed some symptoms of hypoxic such as chlorosis in mature leaves, decreased leave formation, and reduced of root carbohydrates^{13,32}.

The reduction of growth is probably related to reduction of dry matter as informed by Prawoto et al.³⁵, who determined that soil water content 25 % above of field capacity (FC) decreased the dry matter by 13 %, likewise, when the soil water content was 75 % above FC the reduction of dry matter was 34 %. These results are according to other ones informed by de Almeida & Valle¹⁴ and Rehem et al.³². Other traits studied in cocoa cultivars were the relative growth rate (RGR) of leaves and roots which decreased when the plants grew under excessive water^{18,32}. However, the specific leaf area (SLA) increased in the same experiment³².

A probably mechanism to overcome the hypoxic caused by waterlogging is the increment of lenticel on the trunk of cocoa tress as reported by de Almeida et al.¹⁹, Rehem et al.³² and Bertolde et al.³⁶. In this sense, other researchers found overabundance of soil water stimulated the production of hypertrophied lenticels and adventitious roots in *Quercus robur* and *Quercus petraea*³⁷. Other morphological strategy is the formation of adventitious roots³⁶. Furthermore, adaptations of cocoa plants under waterlogging are the reduction of xylem, phloem and mesophyll length³⁸.

Effect on production of cocoa. In spite of there is a lot of information about the effect of waterlogging in different crops, in the case of cocoa this type of information is still scarce. According to de Almeida & Valle¹⁴ the anoxia can reduce the productivity until 60 %.

Strategic adaptation to waterlogging. In order to deal with excessive water on soil, we need to start choosing a vegetative material suitable which be tolerant to flood stress (for instance TSA-792, CP-49)^{32,39}. We must also select a proper production system as agroforestry system (AF). In fact, in an AF the number of survived cocoa plants was higher than cocoa trees unshaded, both growing under flooding⁴⁰. Other strategies recorded were the increment of the glycolysis and morphological modifications (e. g. increase of lenticels)^{36,38} (Figure 3).




FIGURE 3

Strategies to cope with waterlogging for cocoa plantations. To limit the impact of flood on cocoa trees first we have to choose proper production system (for example, agroforestry system). Second, we need to use tolerant cultivar. Internal and external processes such as glycolysis and morphological changes can help cocoa trees to overcome flooding stress

FUTURE PERSPECTIVES

Finally, after searching for information related to the effect of waterlogging on cocoa plants, it has been shown that excess water in the soil (anoxic) can negatively affect the physiology (e. g. reducing net assimilation and stomatal opening), enzymatic activity of peroxidases and polyphenol oxidases, absorption of nutrients (macro and micro nutrients) and the growth of cocoa plants. However, this specie has certain mechanisms to counteract the damage of the lack of oxygen in the root zone, such as the increase of glycolysis. As well, this specie forms lenticels when it is under anoxia. Different agronomic practices can be important to counteract the effect of flood; for example, we have to choose appropriate system of production (in this case agroforestry) and to cultivate flooding tolerant genetic material. In spite of information exists about this issue, it is necessary more research in order to understand better what mechanisms cocoa plants have to survive, or overcome the anoxic condition.

CITED LITERATURE

1. The top cocoa producing countries in the World [Internet]. WorldAtlas. 2021 [citado 5 de junio de 2021]. Recuperado a partir de: <https://www.worldatlas.com/industries/the-top-cocoa-producing-countries-in-the-world.html>

2. Osarenren CO, Ejuetueyin JO, Eweka KI. Socio-economic characteristics of registered cocoa farmers in Edo State, Nigeria. *J Appl Sci Environ Manag* 2016;20(2):261-6. DOI: <https://doi.org/10.4314/jasem.v20i2.5>
3. Hervé BDB, Vidal, S. Plant biodiversity and vegetation structure in traditional cocoa forest gardens in southern Cameroon under different management. *Biodivers Conserv* 2008;17:1821-35. DOI: <https://doi.org/10.1007/s10531-007-9276-1>
4. Ramírez-Meneses A, García-López E, Obrador-Olán JJ, Ruíz-Rosado O, Camacho-Chiu W. Diversidad florística en plantaciones agroforestales de cacao en Cárdenas, Tabasco, México. *Universidad y Ciencia* 2013;29(3):215-30. DOI: <https://doi.org/10.19136/ERA.A29N3.55>
5. Adjaloo M, Oduro W. Insect assemblage and the pollination system of cocoa (*Theobroma cacao* L.). *J Appl Biosci* 2013;62:4582-94. DOI: <https://doi.org/10.4314/jab.v62i0.86070>
6. Tagnang Nadege M, Claude S, Roger K, Cedric C, Ntsomboh N, Yonkeu N, et al. Ecological and economic potentials of cocoa agroforestry systems in the Center Region of Cameroon. *Am J Agric For* 2020;8(5):214-22. DOI: <https://doi.org/10.11648/j.ajaf.20200805.15>
7. Producción nacional de cacao en grano creció en la última década a un promedio de 12.6% al año [Internet]. Ministerio de Desarrollo Agrario y Riego. 2020 [citado el 23 de julio de 2022]. Recuperado a partir de <https://www.gob.pe/institucion/midagri/noticias/305143-produccion-nacional-de-cacao-en-grano-crecio-en-la-ultima-decada-a-un-promedio-de-12-6-al-ano>
8. Niether W, Glawe A, Pfohl K, Adamtey N, Schneider M, Karlovsky P, et al. The effect of short-term vs. long-term soil moisture stress on the physiological response of three cocoa (*Theobroma cacao* L.) cultivars. *Plant Growth Regul* 2020;92: 295-306. DOI: <https://doi.org/10.1007/s10725-020-00638-9>
9. Lahive F, Hadley P, Daymond AJ. The physiological responses of cacao to the environment and the implications for climate change resilience. A review. *Agron Sustain Dev* 2019;39:5. DOI: <https://doi.org/10.1007/s13593-018-0552-0>
10. McCarthy N, Kilic T, Brubaker J, Murray S, Fuente A. Droughts and floods in Malawi: impacts on crop production and the performance of sustainable land management practices under weather extremes. *Environ Dev Econ* 2021;26(5):432-49. DOI: <https://doi.org/10.1017/S1355770X20000455>
11. Yeung E, Bailey-Serres J, Sasidharan R. After the deluge: plant revival post-flooding. *Trends Plant Sci* 2019;24(5):443-54. DOI: <https://doi.org/10.1016/j.tplants.2019.02.007>
12. Anderson R, Bayer PE, Edwards D. Climate change and the need for agricultural adaptation. *Curr Opin Plant Biol* 2020;56:197-202. DOI: <https://doi.org/10.1016/j.pbi.2019.12.006>
13. Bertolde FZ, Almeida AAF, Pirovani CP, Ahnert D, Baligar VC, Valle RR. Physiological and biochemical responses of *Theobroma cacao* L. genotypes to flooding. *Photosynthetica* 2012;50:447-57. DOI: <https://doi.org/10.1007/s11099-012-0052-4>
14. de Almeida AAF, Valle RR. Ecophysiology of the cacao tree. *Braz J Plant Physiol* 2007;19(4):425-48. DOI: <http://doi.org/10.1590/S1677-04202007000400011>
15. Xu Z, Jiang Y, Zhou G. Response and adaptation of photosynthesis, respiration, and antioxidant systems to elevated CO₂ with environmental stress in plants. *Front Plant Sci* 2015;6:701. DOI: <https://doi.org/10.3389/fpls.2015.00701>
16. Plant growth and development [Internet]. Plant Science & Technology. 2022. [citado el 23 de agosto de 2022]. Recuperado a partir de <https://extension.oregonstate.edu/gardening/flowers-shrubs-trees/plant-growth-development>
17. Griffin KL, Seemann JR. Plants, CO₂ and photosynthesis in the 21st century. *Chem Biol* 1996;3 (4):245-254. DOI: [https://doi.org/10.1016/S1074-5521\(96\)90104-0](https://doi.org/10.1016/S1074-5521(96)90104-0)
18. Sena Gomes AR, Kozłowski TT. The effects of flooding on water relations and growth of *Theobroma cacao* var. catongo seedlings. *J Horticult Sci* 1986;61(2):265-76. DOI: <https://doi.org/10.1080/14620316.1986.11515700>
19. de Almeida J, Tezara W, Herrera A. Physiological responses to drought and experimental water deficit and waterlogging of four clones of cacao (*Theobroma cacao* L.) selected for cultivation in Venezuela. *Agric Water Manag* 2016;171:80-8. DOI: <https://doi.org/10.1016/j.agwat.2016.03.012>

20. Rodríguez-Gamir J, Ancillo G, González-Mas MC, Primo-Millo E, Iglesias DJ, Forner-Giner M. Root signalling and modulation of stomatal closure in flooded citrus seedlings. *Plant Physiol Biochem* 2011;49(6):636-45. DOI: <https://doi.org/10.1016/j.plaphy.2011.03.003>
21. De Pedro LF, Mignolli F, Scartazza A, Melana Colavita JP, Bouzo CA, Vidoz ML. Maintenance of photosynthetic capacity in flooded tomato plants with reduced ethylene sensitivity. *Physiol Plant* 2020;170(2):202-17. DOI: <https://doi.org/10.1111/ppl.13141>
22. Fromme P, Jordan P, Krauß N. Structure of photosystem I. *Biochim Biophys Acta Bioenerg* 2001; 1507(1-3):5-31. DOI: [https://doi.org/10.1016/S0005-2728\(01\)00195-5](https://doi.org/10.1016/S0005-2728(01)00195-5)
23. Govindjee JW. Photosystem II. In: *Encyclopedia of Life Sciences*. New York: Macmillan Education; 2002. p. 1-13, DOI: <https://doi.org/10.1038/npg.els.0000669>
24. González Moreno S, Perales Vela H, Salcedo Álvarez MO. La fluorescencia de la clorofila a como herramienta en la investigación de efectos tóxicos en el aparato fotosintético de plantas y algas. *REB* 2008;27(4):119-29.
25. Costa RJ, Ferreira TR, Silva FR e, Souza GAR de, Cerri Neto B, Leite L de A, et al. Fluorescência transiente da clorofila a em plantas de *Theobroma cacao* submetidas a diferentes períodos de alagamento. In: *Encontro Latino-Americano de Iniciação Científica, 22.; Encontro Latino Americano de Pós Graduação, 18.; Encontro Nacional de Iniciação À Docência, 8, 2017, Urbanova. Patrimônio e Cultura Desafios da Ciência Frente a Identidades Plurais [Internet]. São José dos Campos: UNIVAP; 2018 [citado 3 de mayo de 2020]. p. 1-4. Recuperado a partir de: <https://biblioteca.incaper.es.gov.br/digital/bitstream/123456789/3569/1/RE-1065-1255-08.pdf>*
26. Christen D, Schönmann S, Jermini M, Strasser RJ, Défago G. Characterization and early detection of grapevine (*Vitis vinifera*) stress responses to esca disease by *in situ* chlorophyll fluorescence and comparison with drought stress. *Environ Exp Bot* 2007;60(3):504-14. DOI: <https://doi.org/10.1016/j.envexpbot.2007.02.003>
27. Živčák M, Olšovská K, Slamka P, Galambošová J, Rataj V, Shao H, et al. Application of chlorophyll fluorescence performance indices to assess the wheat photosynthetic function influenced by nitrogen deficiency. *Plant Soil Environ* 2014; 60(5):210-15. DOI: <https://doi.org/10.17221/73/2014-PSE>
28. Swoczyna T, Lata B, Stasiak A, Stefaniak J, Latocha P. JIP-test in assessing sensitivity to nitrogen deficiency in two cultivars of *Actinidia arguta* (Siebold et Zucc.) Planch. ex Miq. *Photosynthetica* 2019;57(2):646-58. DOI: <https://doi.org/10.32615/ps.2019.057>
29. da Silva Branco MC, de Almeida AAF, Dalmolin AC, Ahnert D, Baligar VC. Influence of low light intensity and soil flooding on cacao physiology. *Sci Hortic* 2017;217:243-57. DOI: <https://doi.org/10.1016/j.scienta.2017.01.038>
30. de Souza Oliveira V, Braidão Pinheiro AP, Cerri Neto B, Dousseau Arantes S, Alves da Silva C, Crasque J. Effect of flooding under the gas exchange of cocoa seedlings. *J Agric Sci* 2019; 11(16):233-7. DOI: <https://doi.org/10.5539/jas.v11n16p233>
31. Perata P, Alpi A. Plant responses to anaerobiosis. *Plant Sci* 1993;93(1-2):1-17. DOI: [https://doi.org/10.1016/0168-9452\(93\)90029-Y](https://doi.org/10.1016/0168-9452(93)90029-Y)
32. Rehem BC, de Almeida AAF, Mielke MS, Gomes FP. Effects of substrate flooding on growth and chemical composition of *Theobroma cacao* L. clonal genotypes. *Rev Bras Frutic* 2009;31(3): 805-15. DOI: <https://doi.org/10.1590/S0100-29452009000300026>
33. Rasheed R, Iqbal M, Arslan Ashraf M, Hussain I, Shafiq F, Yousaf A, et al. 2018. Glycine betaine counteracts the inhibitory effects of waterlogging on growth, photosynthetic pigments, oxidative defence system, nutrient composition, and fruit quality in tomato. *J Hortic Sci Biotechnol* 2018;93(4): 385-91. DOI: <https://doi.org/10.1080/14620316.2017.1373037>
34. Wollmer AC, Pitann B, Mühling KH. Waterlogging events during stem elongation or flowering affect yield of oilseed rape (*Brassica napus* L.) but not seed quality. *J Agron Crop Sci* 2018; 204(2):165-74. DOI: <https://doi.org/10.1111/jac.12244>
35. Prawoto A, Zainunnuroni M, S. Response of selected clones of cocoa seedlings in the nursery against high soil water content. *Pelita Perkebunan* 2005;21(2):90-105. DOI: <https://doi.org/10.22302/iccri.jur.pelitaperkebulan.v21i2.17>

36. Bertolde FZ, De Almeida AAF, Corrêa RX, Gomes FP, Gaiotto FA, Baligar VC, et al. Molecular, physiological and morphological analysis of waterlogging tolerance in clonal genotypes of *Theobroma cacao* L. *Tree Physiology* 2010; 30(1):56-67. DOI: <https://doi.org/10.1093/treephys/tpp101>
37. Le Provost G, Lesur I, Lalanne C, Da Silva C, Labadie K, Aury JM, et al. Implication of the suberin pathway in adaptation to waterlogging and hypertrophied lenticels formation in pedunculate oak (*Quercus robur* L.). *Tree Physiol* 2016;36(11): 1330-42. DOI: <https://doi.org/10.1093/treephys/tpw056>
38. Silva FR de, Souza GAR de, Ferreira T, Costa RJ, Correia LZ, Neto B, Souza CAS. Caracterização anatômica e pigmentos fotossintéticos de folhas de mudas de cacaueteiro submetidas ao alagamento. In: XII Encontro Latino Americano de Iniciação Científica Júnior da Univap [Internet]. Universidad do Vale do Paraíba: UNIVAP; 2018 [citado 3 de mayo de 2020]. p. 1-6. Recuperado a partir de: http://www.inicepg.univap.br/cd/INIC_2018/anais/arquivos/RE_0238_0706_01.pdf
39. Bertolde FZ, Almeida AA, Pirovani CP. Analysis of gene expression and proteomic profiles of clonal genotypes from *Theobroma cacao* subjected to soil flooding. *PLoS One* 2014;9(10): e108705. DOI: <https://doi.org/10.1371/journal.pone.0108705>
40. Delgado C, Penn J, Couturier G. Status of cocoa trees following seasonal floods in major watersheds of the Peruvian Amazon. *Agric Sci* 2016; 4(2):15-24. DOI: <https://doi.org/10.12735/as.v4n2p15>

NOTES

Funding source: This work did not receive funding.

Conflicts of interest: The authors state there is no conflicts of interest.

Acknowledgements: We want to thank to Universidad Nacional Agraria La Molina.

Ethical considerations: The authors declare there is no plagiarism in this manuscript.

Authors' contributions: Saravia-Castillo Gabriela, Castro-Cepero Viviana, Julca Otiniano Alberto, Alvarado-Huamán Leonel, Borjas-Ventura Ricardo, searched, organized and analyzed the information and reviewed the final version of this document.

Article ID: 117/JSAB/2022

Editor's Note: *Journal of the Selva Andina Biosphere (JSAB)*. All statements expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, editors and reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

ALTERNATIVE LINK

http://www.scielo.org/bo/scielo.php?script=sci_arttext&pid=S2308-38592022000200078&lng=es&nrm=iso&tlng=en (html)